



可視光サーベイデータの 使い方・見方 ~PanSTARRS1を中心に~




小久保 充 (東北大学)

Contents

- インTRODクシヨN: SDSSのデータを見る
- 可視撮像データの基礎概念 (分光は扱わない)
- PanSTARRS1パブリックデータリリース
 - 画像, 測光カタログ
 - データベースの利用
- 他のパブリックサーベイデータ



Sloan Digital Sky Survey



SDSS-I 2000-2005
SDSS-II 2005-2008
SDSS-III 2008-2014
SDSS-IV 2014-2020
SDSS-V 2020 —

撮像は2000-2001で完了
→ 撮像選択された天体を分光

Band	5 σ	Saturation
<i>u</i>	22.15	13
<i>g</i>	23.13	14
<i>r</i>	22.70	14
<i>i</i>	22.20	14
<i>z</i>	20.71	12




大きなデジカメ

Sloan Digital Sky Survey





データベースに保管されている各画像は、
Filter
Run
Field
Camera Column
解析version(rerun)
で指定される。



driftscan (run)
(各sky位置が54秒で1CCDを横切る)

Sloan Digital Sky Survey

最新はData Release (DR) 16




- SDSS DR16 Navigator Tool
 - 座標移動, zoom-in, drawing options
 - Explore
 - photoObj table
 - Fits画像データ取得

<http://skyserver.sdss.org/dr16/en/tools/chart/navi.aspx>


<http://skyserver.sdss.org/dr16/en/tools/explore/Summary.aspx?id=1237663783138361393>

安田さんによる資料: https://www.adc.nao.ac.jp/people/~furusawa/work/da_ws/2/slides/Yasuda_SDSS.pdf

SDSS測光値の定義: <https://www.sdss.org/dr12/algorithms/magnitudes/>



Sloan Digital Sky Survey : 画像データ



- 54秒積分 (1 epochのみ; 2000-2001)
- サーベイ領域の**モザイク画像**が
2k × 1k 画像(*frame*)に小分けされている.
FITS形式.
- Sky引き算, WCS座標校正済み
- 物理的に意味のあるcountに校正済み:
ゼロ点等級 = 22.5 mag (1 nanomaggy
[1 count が 22.5 mag に対応])
- カタログの測光値を使う or 自分で画像の測光を行う
(銀河などのextended sourceは自分で画像を確認した
ほうが良いだろう)

<http://dr16.sdss.org/sas/dr16/eboss/photoObj/frames/301/4263/2/frame-g-004263-2-0340.fits.bz2>


https://www.sdss.org/dr16/imaging/imaging_basics/

<https://www.sdss.org/dr16/algorithms/magnitudes/#nmgy>



Python access (astroquery) : <https://astroquery.readthedocs.io/en/latest/sdss/sdss.html>

Sloan Digital Sky Survey : 画像データ

カウント
合計






等級



- 星のカatalog測光値としては**PSF等級**(PSF profileをfitしたもの; psfMag in *PhotoObj* table)を使う.
- 十分大きな開口をとってカウントを合計したものがPSF等級と等価.
(小さい開口をとる場合はaperture correctionが必要)

Sloan Digital Sky Survey : 画像データ




- 広がった天体の測光値は色々な定義がある
- modelMag [in *PhotoObj* table] がSDSSの推奨.
(DeV, Exp, ... モデルプロファイルfitによる測光値)
- ペトロシアン半径の開口測光値なども利用できる
(in *PhotoObj* table)
- ds9でquickにでも自分で測光してみるのがbest



Sloan Digital Sky Survey : 画像データ



Image



$\text{sigma_map} = \text{sqrt}(\text{variance map})$



gain = 3.855 (tsField fileから取ってくる) : https://dr16.sdss.org/datamodel/files/SDSS_TARGET/data/tsField.html

Background RMSは photutils.background で見積もった: <https://photutils.readthedocs.io/en/stable/background.html>

Sloan Digital Sky Survey : 測光カタログ

一般的なカタログ (tables) アクセスはSQL : / SQL queries 実用例



The screenshot shows the SkyServer DR16 plus web interface. The top navigation bar includes links for Home, Data, Schema, Education, Astronomy, SDSS, Contact Us, Download, Site Search, Help, and History (marked as NEW!). The left sidebar lists various tools under 'DR16 Tools', including 'Getting Started', 'Famous places', 'Get images', 'Scrolling sky', 'Visual Tools', 'Search' (with sub-options: Radial, Rectangular, Search Form, SQL, Imaging Query, Spectro Query, IR Spec Query), 'CrossID', 'Skyquery CrossMatch', and 'CasJobs'. The main content area is titled 'SQL Search' and contains a text box for entering SQL queries. A sample query is provided, which joins the PhotoObj and SpecObj tables and selects the top 10 results based on magnitude and color. Below the query box are buttons for 'Check syntax', 'Submit query', and 'Clear Query'. There is also a section for 'Output Format' with radio buttons for HTML (selected), XML, CSV, JSON, VOTable, FITS, and MyDB (marked as NEW!). A 'Table name' input field and a 'Reset' button are also present. At the bottom, there are links to the 'Schema Browser' and a 'Searching for Data' tutorial, along with a detailed explanation of the query requirements for data upload.

SLOAN DIGITAL SKY SURVEY
SkyServer DR16 plus

Home Data Schema Education Astronomy SDSS Contact Us Download Site Search Help History **NEW!**

DR16 Tools

- Getting Started
- Famous places
- Get images
- Scrolling sky
- Visual Tools
- Search
 - Radial
 - Rectangular
 - Search Form
 - **SQL**
 - Imaging Query
 - Spectro Query
 - IR Spec Query
- CrossID
- Skyquery CrossMatch
- CasJobs

SQL Search

This page allows you to directly submit a [SQL \(Structured Query Language\)](#) query to the SDSS database server. You can modify the default query as you wish, or cut and paste a query from the [SDSS Sample Queries page](#).

Please note: To be fair to other users, queries run from SkyServer search tools are restricted in how long they can run and how much output they return, by **timeouts** and **row limits**. Please see the [Query Limits help page](#). To run a query that is not restricted by a timeout or number of rows returned, please use the [CasJobs batch query service](#).

```
-- This query does a table JOIN between the imaging (PhotoObj) and spectra
-- (SpecObj) tables and includes the necessary columns in the SELECT to upload
-- the results to the SAS (Science Archive Server) for FITS file retrieval.
SELECT TOP 10
  p.objid,p.ra,p.dec,p.u,p.g,p.r,p.i,p.z,
  p.run, p.rerun, p.camcol, p.field,
  s.specobjid, s.class, s.z as redshift,
  s.plate, s.mjd, s.fiberid
FROM PhotoObj AS p
JOIN SpecObj AS s ON s.bestobjid = p.objid
WHERE
  p.u BETWEEN 0 AND 19.6
  AND g BETWEEN 0 AND 20
```

Output Format ☒ HTML ☐ XML ☐ CSV ☐ JSON ☐ VOTable ☐ FITS ☐ MyDB **NEW!**

Table name

To find out more about the database schema use the [Schema Browser](#).

For an introduction to the Structured Query Language (SQL), please see the [Searching for Data](#) How-To tutorial. In particular, please read the [Optimizing Queries](#) section.

The inclusion of the imaging and spectro columns for [SAS](#) upload in your query (as in the default query on this page) will ensure that when you press **Submit**, the appropriate button(s) are displayed on the query results page to allow you to upload the necessary information to the [SAS](#) to retrieve the FITS file data corresponding to your CAS query. The imaging columns needed for upload to the [SAS](#) are *run*, *rerun*, *camcol*, and *field*. The spectroscopic columns needed are *plate*, *mjd*, *fiberid*, and optionally *sprerun* (the latter requires a join with the PlateX table).

- 測光カタログと分光カタログをくっつける。
- 条件にあう天体を10個表示

SDSSのSQL web interface: <http://skyserver.sdss.org/dr16/en/tools/search/sql.aspx>

Sloan Digital Sky Survey : 測光カタログ

SQLへのpython access

大量データ取得や,
変動天体探査自動pipelineへの
組み込み等で有用

SQL query定義

```
: import sqlcl
import numpy as np
import pandas as pd
```

```
: # http://skyserver.sdss.org/dr16/en/tools/search/sql.aspx
```

```
sql_script="\n
SELECT TOP 10 \n
    p.objid,p.ra,p.dec,p.u,p.g,p.r,p.i,p.z, \n
    p.run, p.rerun, p.camcol, p.field, \n
    s.specobjid, s.class, s.z as redshift, \n
    s.plate, s.mjd, s.fiberid \n
FROM PhotoObj AS p \n
    JOIN SpecObj AS s ON s.bestobjid = p.objid \n
WHERE \n
    p.u BETWEEN 0 AND 19.6 \n
    AND g BETWEEN 0 AND 20 \n
"
```

SQL query submit

```
: lines = sqlcl.query(sql_script).readlines()
print(lines)

data = []
for line in lines[2:]:
    #data.append(map(float,line[:-1].split(',')))
    data.append( line[:-1].split(','))

columns=lines[1].split(",")

catalog = pd.DataFrame( data, columns=columns )
```

任意のtable形式に変換

	objid	ra	dec	u	g	r	i	z	run	rerun	camcol	field	spec
0	1237645879551066262	348.902530171573	1.27188624932946	19.38905	18.24496	17.58728	17.20807	16.90905	94	301	6	94	4301949499510
1	1237645879578460255	51.4436950026148	1.27007271795861	19.52808	17.96541	17.03493	16.53754	16.14154	94	301	6	512	4662352927313
2	1237645879578460271	51.4835835170322	1.27201267342336	18.72268	17.3852	16.81134	16.51803	16.29502	94	301	6	512	4662328188301
3	1237645941291614227	49.6274851210218	-1.0417691497987	17.65612	16.17133	15.5894	15.3785	15.26744	109	301	1	100	17057955826620
4	1237645941824356443	40.2856904119424	-0.71495662825756	17.54884	15.75164	15.031	14.66728	14.36099	109	301	2	37	4593946803338
5	1237645941824356481	40.2721052364832	-0.642510265945889	19.23401	17.5333	16.8743	16.63157	16.49762	109	301	2	37	12047442798727
6	1237645942898229350	40.5820323404104	0.134770090982145	18.64558	16.44336	15.52452	15.18185	14.98858	109	301	4	39	82531204825699
7	1237645942905438473	57.025336578067	0.20884499634774	17.61444	16.17125	15.52131	15.15564	14.86996	109	301	4	149	13984884043704
8	1237645942905438495	57.0470524254038	0.0843042629762005	19.46874	18.18264	17.59063	17.26436	16.95295	109	301	4	149	17216016120335
9	1237645942905569371	57.281615211522	0.0187678802337251	16.4848	14.92993	14.56054	14.53054	14.19394	109	301	4	151	13984985748529

sqlcl: <http://skyserver.sdss.org/dr16/en/help/download/sqlcl/sqlcl.aspx>

@Haleakala (3040m), Hawaii

北天

PanSTARRS1 3π survey (完了)

DR2 (2019-01-28), Chambers+2016

g,r,i,z,y-band (4000-10000 Å)

Equatorial



i_{P1} - band magnitude for 98% completeness

@Siding Spring
Observatory (1165m),
Australia.


南天


1.35m

u,v,g,r,i,z-band (3500-10000 Å)


SkyMapper Southern Survey (進行中)



DR2 (2019-02-27), Onken+2019





Deeper (=narrower) imaging surveys






(← image測光を自分でやり直した)

多くの可視・赤外線レガシーサーベイデータが自由に利用できる
→ 特にtime-domainで有用


可視撮像データの基礎

望遠鏡, フィルター, CCD検出器

光 大気



光




光

高い線型性




光電子
→ デジタル化





Sky Poisson noise
(+read noise)

Object



モザイクCCD



- 望遠鏡焦点面の視野を埋め尽くすように複数のCCD chipを並べる
→ モザイクCCDカメラ (SDSS, PanSTARRS, SkyMapper, HSC, VISTA, LSST, ...)
- chip間に物理的隙間ができる
→ 少しずつ視野をずらして **足し算** (coadd, stack) することで
隙間を埋める+**限界等級を深くする**



Subaru/Hyper Suprime-Cam
116 CCD chips

PanSTARRS1/GPC1

60 CCD chips



AB等級

- Response関数 (=フィルター透過 × 光学系 × CCD量子効率…) × 光子数重みづけしたfluxが, そのフィルターでの <flux密度> :



$$f_\nu \leftarrow \frac{\int R_\nu \times \frac{f_\nu}{\nu} d\nu}{\int R_\nu \times \frac{1}{\nu} d\nu}$$

- AB等級: 3631 Jansky (3.631×10^{-20} cgs) の星を0等星とする
現代の可視光サーベイの測光システム”ほぼ”AB等級
(実際は~1%くらいズれているので、1%精度の話をしたときは注意)


$$\begin{aligned} m_{AB}(\nu) &= -2.5 \log(f_\nu / 3631 \text{ Jy}) \\ &= -48.600 - 2.5 \log(f_\nu [\text{erg/sec/cm}^2/\text{Hz}]) \end{aligned}$$

$$\Delta m_{AB} = -1.086 \times \frac{\Delta f_\nu}{f_\nu} \sim \frac{\Delta f_\nu}{f_\nu}$$


e.g., 10%のフラックス光度変動 ~ 0.1magの光度変動
1%のフラックス精度 ~ 0.01magの精度



Wide-field imaging surveys



撮像深さ
(限界等級)



天域の広さ

- そこそこの口径の望遠鏡で超広視野 → SDSS, PanSTARRS, SkyMapper
- 大口径の望遠鏡でそこそこの広視野 → Blanco/DECam, Subaru/HSC, (LSST)

CCD image

- Pixelカウントの単位はADU (Analog-Digital Unit):

$$\text{photo-electrons}[e^-] = \text{gain}[e^-/\text{ADU}] \times \text{counts}[\text{ADU}]$$

$$\text{Poisson error}[e^-] = \sqrt{\text{photo-electrons}}$$

$$\begin{aligned} \text{Poisson error}[\text{ADU}] &= \sqrt{\text{photo-electrons}} / \text{gain} \\ &= \sqrt{\text{counts} / \text{gain}} \end{aligned}$$

where **gain** $[e^-/\text{ADU}]$ is an instrument-specific constant (readout electronics)

- Pixelカウント


$$= (\text{Bias} + \text{Dark current} + \text{Sky background} + \text{Cosmic rays} + \text{Objects})$$

$$\times \text{Sensitivity variation} + \text{random noise}$$

1次処理後,


$$\text{Pixelカウント} = \text{Objects} + \text{random noise}$$

- 等級ゼロ点較正, WCS座標較正など.




PanSTARRS1 60 CCDs; Waters+2016

Resampled *warp* と *stack* images



PanSTARRS 3 π RINGS.V3 tessellations; Chambers+2016

- 1次処理した画像を, 天球面を平面に分割した **pre-definedセル**上に投影する.
 - 画像の回転, ピクセルresamplingが必要 ... *warp*
(画像のpixelサイズはあまり変えないようにする; PanSTARRSでは $\sim 0.26''/\text{pix} \rightarrow 0.25''/\text{pix}$)
- 複数の **single-epoch warp**画像を足して (*stack*) して, 隙間のない deepなモザイク画像をつくる





warp



PanSTARRS1 60 CCDs; Waters+2016

分散 Variance map



- 自分でstack画像の測光を行う場合, 測光値のエラーを正しく推定するのは結構難しい.

σ_{sky} = Sky光 Poisson (+ readout) noise

σ_{obj} = Object光 Poisson noise

Total variance = $(\sigma_{\text{sky}})^2 + (\sigma_{\text{obj}})^2$

- Stackingの過程で誤差伝播した最終variance [各pixelのGaussian variance σ^2] の画像 = Variance map を提供しているサーベイもある (e.g., PanSTARRS1)
→ variance mapに対して同じ開口で”測光”して、 $\sqrt{\text{variance_sum}}$ でエラー評価




$$\begin{aligned} eFlux(1\sigma) &= \sqrt{\sum_{\text{pixel}} Variance_{\text{pixel}}} \\ &= \sqrt{\sum_{\text{pixel}} \sigma_{\text{pixel}}^2} \end{aligned}$$

Waters+2016


Mask image

- 飽和した明るい星, 宇宙線, chip gapなどにより使えない pixelの画像 (e.g., good pixel=0, bad pixel=1)
- Stackingすることで, maskされる領域は減る


Waters+2016




Warp image



Mask image



Stack image



Mask image


Data Release 2

PanSTARRS1 DR2 public data

- 画像: warp画像, stack画像(+mask画像,variance画像,...)
- 測光カタログ: warp測光, stack測光, warp forced測光

PanSTARRS 3 π ; g, r, i 3色合成


Chambers+2016





Pan-STARRS



- Panoramic Survey Telescope and Rapid response System (PanSTARRS; PS)
- もともとの計画は4台の同型の**1.8m口径 telescopes** → budget problem
 - **PS1 & PS2** telescopes constructed
 - **Public survey was carried out by the PS1 telescope in 2010-2014.**
- Wide-field time-domain surveys (Near Earth Objects, Asteroids, Galactic variables, SNe ...)
- Deep stack images for extragalactic studies



Giga Pixel Camera 1 (GPC1)



- 1.4 Gigapixels (14億ピクセル/1露出) mosaic camera
- 特に長波長側で高感度



The PS1 Surveys

Survey	Area (sq.deg.)	Filters	Percent Time	Exposure (secs)
3 π	30,000	g,r,i,z,y	56	43/40/45/30/30
Medium Deep	70	g,r,i,z,y	25	113/113/240/240/ 240
Solar System	10,000	w	5	45
Stellar Transit	40	i	4	15 or 30
M31	7	r,i	2	60/60

DR2

DR3?


Data collection started early in 2010 and completed in April 2014

PS1 3π -steradian survey

- 2010 — 2014 (3.5 yrs), *g, r, i, z, y*-band
- 3π 北天scan (Decl. $< -30^\circ$):
各バンド12 exposures over 3.5 yrs
- **22-23 mag (final stack).** SDSSより~1 – 2等深い
- Excellent seeing (~1 arcsec in median)


Detection limits (AB mag.)

Band	5 σ (single)	5 σ (stack)	Saturation
<i>g</i>	22.0	23.3	14.5
<i>r</i>	21.8	23.2	15.0
<i>i</i>	21.5	23.1	15.0
<i>z</i>	20.9	22.3	14.0
<i>y</i>	19.7	21.4	13.0



PS1 Data Release 2 (DR2) data product

- Pipeline(PV3)-reduced data products are available at STScI/MAST:
 - **Single-epoch warp images**
 - **Stack images**
 - **たくさんのphotometry catalogs**
- **Not** included in the release (to be released in DR3):
 - Subtracted images and catalogues for transient detections
 - Data from other surveys (Medium Deep survey ...)



<https://star.pst.qub.ac.uk/ps1threepi/psdb/>

IMAGE PROCESSING PIPELINE STAGES AND DATA PRODUCTS

processing stage	image class	file type	ID type	Avg No. components per ID	No. filters per ID	No. IDs in 3π Survey	location	Release
raw	raw image	fits	exp	60.0000	1	374k	UH ^a	...
chip	signal image	fits	exp	59.9984	1	374k	... ^b	...
	variance image	fits	exp	59.9984	1	374k	...	...
	mask image	fits	exp	59.9984	1	374k	...	...
	detections	cmf	exp	59.9984	1	374k	both ^c	...
camera	detection table	smf	exp	1	1	374k	both	...
warp	signal image	fits	skycell	72.6060	1	1050k	MAST	DR2
	variance image	fits	skycell	72.6060	1	1050k	MAST	DR2
	mask image	fits	skycell	72.6060	1	1050k	MAST	DR2
	detections	cmf	skycell	72.6060	1	1050k	both	...
stack	signal image	fits	skycell	1	1	1050k	both	DR1
	variance image	fits	skycell	1	1	1050k	both	DR1
	mask image	fits	skycell	1	1	1050k	both	DR1
	number image	fits	skycell	1	1	1050k	both	DR1
	exp image	fits	skycell	1	1	1050k	both	DR1
	expwt image	fits	skycell	1	1	1050k	both	DR1
	convolved 6&8 signal	fits	skycell	1	1	1050k	...	...
	convolved 6&8 variance	fits	skycell	1	1	1050k	...	...
	convolved 6&8 mask	fits	skycell	1	1	1050k	...	...
	convolved 6&8 number	fits	skycell	1	1	1050k	...	...
	convolved 6&8 exp	fits	skycell	1	1	1050k	...	...
	convolved 6&8 expwt	fits	skycell	1	1	1050k	...	...
	detections	cmf	skycell	1	1	1050k	both	...
static sky	detections	cmf	skycell	1	5	201k	both	...
sky cal	detections	cmf	skycell	1	1	1050k	both	...
forced	detections	cmf	forced	19.1184	1	19m	both	...
diff	signal image	fits	skycell	51.6105	1	19m	...	...
	variance image	fits	skycell	51.6105	1	19m	...	...
	mask image	fits	skycell	51.6105	1	19m	...	...
	diff detections	cmf	skycell	51.6105	1	19m	both	DR2




FUNDAMENTAL PSPS DATABASE TABLES

Table Class	PSPS Table Name	Release		
Detection	Detection	DR2		
Object	ObjectThin	DR1		
	MeanObject	DR1		
Stack	GaiaFrameCoordinate	DR1	3π STERADIAN SURVEY PHOTOMETRY SCOPE	
	StackObjectThin	DR1		
	StackObjectAttributes	DR1		
	StackApFlx	DR1	Measurement	Scope
	StackApFlxExGalUnc	DR1	PSFMag	All objects
	StackApFlxExGalCon6	DR1	ApMag	All objects
	StackApFlxExGalCon8	DR1	KronMag	All objects
	StackPetrosian	DR1	Fixed apertures R5-R7	All objects
	StackModelFitExp	DR1	Fixed apertures R3-R4, R8-R11	Objects outside Galactic plane
	StackModelFitDeV	DR1	Petrosian	Objects outside Galactic plane with S/N > 20
Difference	StackModelFitSer	DR1	Sersic	Objects outside Galactic plane with S/N > 20
	DiffDetection	DR2	Exponential	Objects outside Galactic plane with S/N > 20
	DiffDetObject	DR2	de Vaucouleurs	Objects outside Galactic plane with S/N > 20
Forced	ForcedMeanObject	DR1		
	ForcedWarpMeasurement	DR2		
	ForcedMeanLensing	DR2		
	ForcedWarpLensing	DR2		
	ForcedGalaxyShape	DR2		
	ForcedWarpExtended	DR2		
	ForcedWarpMasked	DR2		

TABLE 9
CURRENTLY DEFINED VIEWS WITHIN PSPS


View Name	Tables used to create view
DetectionObjectView	ObjectThin, MeanObject, Detection
MeanObjectView	ObjectThin, MeanObject
StackObjectView	ObjectThin, StackObjectThin, StackObjectAttributes
StackApFlxObjectView	ObjectThin, StackApFlx
StackApFlxExGalUncObjectView	ObjectThin, StackApFlxExGalUnc
StackApFlxExGalCon6ObjectView	ObjectThin, StackApFlxExGalCon6
StackApFlxExGalCon8ObjectView	ObjectThin, StackApFlxExGalCon8
StackModelObjectView	ObjectThin, StackModelFitExp, StackModelFitDeV, StackModelFitSer, StackPetrosian
StackModelFitExpObjectView	ObjectThin, StackModelFitExp
StackModelFitDeVObjectView	ObjectThin, StackModelFitDeV
StackModelFitSerObjectView	ObjectThin, StackModelFitSer
StackModelFitPetObjectView	ObjectThin, StackPetrosian
StackObjectPrimaryView	ObjectThin, StackObjectThin, StackObjectAttributes, StackApFlx
DiffDetObjectView	DiffDetObject, DiffDetection
ForcedDetObjectView	ObjectThin, ForcedWarpMeasurement
ForcedMeanObjectView	ObjectThin, ForcedMeanObject
ForcedGalaxyModelView	ObjectThin, ForcedGalaxyShape

Sky Viewer (AladinLite)




Skycells in RINGS.V3 tessellations

- **skycell** : PanSTARRS1 imageの最小解析単位
 - PanSTARRS1 の *warp* 投影座標の定義.
 - 全天 = 2643個のprojection cell, each projection cell = 10×10 **skycells**
 - **skycell** = $25' \times 25'$ (6000×6000 pixel)




Data product: skycell images

skycell.1240.080 (6000 × 6000 pixels = 25' × 25')




Warp (single-epoch; 43sec)

(大きい隙間はCCD gap,
細かい四角はreadout gap)



Warp variance map (wt)



Warp mask

<https://ps1images.stsci.edu/cgi-bin/fitcut.cgi>で取得 (後述)

Warp
Warp.wt
Warp.mask


https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.wt.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.wt.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits

Data product: skycell images

skycell.1240.080 (6000 × 6000 pixels = 25' × 25')



← CCD gap以外に
OTA (Orthogonal Transfer Array)のreadoutによる
隙間が多くある. warpの開口測光は難しい(profile fitに頼る)

Warp (single-epoch; 43sec)

(大きい隙間はCCD gap,
細かい四角はreadout gap)


Warp
Warp.wt
Warp.mask

<https://ps1images.stsci.edu/cgi-bin/fitcut.cgi>で取得 (後述)

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.fits
https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.wt.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.wt.fits
https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits

Data product: skycell images

skycell.1240.080 (6000 × 6000 pixels = 25' × 25')



Warp (single-epoch; 43sec)

(大きい隙間はCCD gap,
細かい四角はreadout gap)

```
python  
from astropy.io import fits
```

```
hdu = fits.open("cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.fits")
```

```
print( "GAIN,SEEING,MAG_ZP,EXPTIME", hdu[0].header['HIERARCH CELL.GAIN'],  
hdu[0].header['HIERARCH CHIP.SEEING']  
, hdu[0].header['HIERARCH FPA.ZP']  
, hdu[0].header['EXPTIME'])
```

```
( 'GAIN,SEEING,MAG_ZP,EXPTIME', 0.9995898, 5.494817, 24.49128, 43.0)
```

<https://ps1images.stsci.edu/cgi-bin/fitcut.cgi>で取得 (後述)

Warp
Warp.wt
Warp.mask

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.wt.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.wt.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits


https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits


https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.wrp.g.55479_49916.mask.fits

Data product: skycell images


skycell.1240.080 (6000 × 6000 pixels = 25' × 25')



Stack(~400sec)



Stack variance map (wt)



Stack mask

<https://ps1images.stsci.edu/cgi-bin/fitcut.cgi>で取得 (後述)

Stack

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.stk.g.unconv.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.stk.g.unconv.fits


Stack.wt

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.stk.g.unconv.wt.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.stk.g.unconv.wt.fits


Stack.mask

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.stk.g.unconv.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.stk.g.unconv.mask.fits

Data product: skycell images



Stack(~400sec)



Mrk590

(等級ゼロ点は 25.0等になっている)

<https://ps1images.stsci.edu/cgi-bin/fitcut.cgi>で取得 (後述)

Stack

Stack.wt

Stack.mask

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.stk.g.unconv.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.stk.g.unconv.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.stk.g.unconv.wt.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.stk.g.unconv.wt.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.stk.g.unconv.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.stk.g.unconv.mask.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.stk.g.unconv.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.stk.g.unconv.mask.fits

https://ps1images.stsci.edu/cgi-bin/fitcut.cgi?red=/rings.v3.skycell/1240/080/rings.v3.skycell.1240.080.stk.g.unconv.mask.fits&format=fits&x=33.639840&y=-0.766690&size=18000&wcs=1&imagename=cutout_rings.v3.skycell.1240.080.stk.g.unconv.mask.fits

Data product: skycell images

- Web interface (PanSTARRS-1 Image Access)で視覚的にcheck & downloadできる



PanSTARRS-1 Image Access

MRK590 [Help](#)

Filters: ☒ color ☒ g ☒ r ☐ i ☐ z ☐ y

File types: ☒ stack ☒ warp


Auxiliary data: ☒ data ☒ mask ☒ wt ☒ exp ☒ expwt ☒ num

Cutout image size: pixels (50.00 arcsec) (sets spatial size of the FITS image)

JPEG display size: pixels (sets resolution of the JPEG previews)

MRK590 (ra = 33.639840, dec = -0.766690)


stack 1240.080 r/g
[Display](#)



stack 1240.080 g
[Display](#) [FITS](#) [FITS-cutout](#)



stack 1240.080 r
[Display](#) [FITS](#) [FITS-cutout](#)



- data: stack, warp images
- mask: mask image
- weight: variance
- exp: 各pixelのexposure timeのmap
- num: 各pixelにstackされたwarpの枚数

warp 1240.080 g 55479.41146
[Display](#) [FITS](#) [FITS-cutout](#)



warp 1240.080 g 55479.49916
[Display](#) [FITS](#) [FITS-cutout](#)



warp 1240.080 g 55508.33769
[Display](#) [FITS](#) [FITS-cutout](#)



warp 1240.080 g 55508.34881
[Display](#) [FITS](#) [FITS-cutout](#)



warp 1240.080 g 55863.38367
[Display](#) [FITS](#) [FITS-cutout](#)



warp 1240.080 g 55863.39528
[Display](#) [FITS](#) [FITS-cutout](#)



<https://ps1images.stsci.edu/cgi-bin/ps1cutouts>

詳細, 注意点(必読) → <https://outerspace.stsci.edu/display/PANSTARRS/PS1+Image+Cutout+Service>

Data product: Photometry Catalogues

主なカタログを理解するための用語:

- Warp imageでの検出 (**Detections**)
 - 1秒角以内のDetectionsを **Object** としてグループ化
 - 各Objectsの複数epochの平均値 (**MeanObject**)
- Stack画像上の**Object** (**StackObject**)
 - … 暗い天体まで検出できるが, stackingによる系統誤差(PSFの形).
- Stack画像上の**Object**の検出位置・形状パラメータをinputとして、warp画像上で**ForcedWarp**測光
 - … warp画像単体では検出できていない暗い天体の測光値を得る
- 上記**ForcedWarp**測光の平均値をとったもの (**ForcedMean**)
 - … StackObjectsよりも系統誤差の少ない高精度測光ができる

測光テーブル


Table Class	PSPS Table Name	Release
Detection	Detection	DR2
	Object	DR1
Stack	MeanObject	DR1
	GaiaFrameCoordinate	DR1
	StackObjectThin	DR1
	StackObjectAttributes	DR1
	StackApFlx	DR1
	StackApFlxExGalUnc	DR1
	StackApFlxExGalCon6	DR1
	StackApFlxExGalCon8	DR1
	StackPetrosian	DR1
	StackModelFitExp	DR1
	StackModelFitDeV	DR1
	StackModelFitSer	DR1
	Forced	DR1
	ForcedMeanObject	DR1
Forced	ForcedWarpMeasurement	DR2
	ForcedMeanLensing	DR2
	ForcedWarpLensing	DR2
	ForcedGalaxyShape	DR2
	ForcedWarpExtended	DR2
	ForcedWarpMasked	DR2

Chambers+2016, Flewelling+2016

$g-i$ VS. i color-magnitude for stars in Messier 53

- **MeanObject** カタログは single exposureの深さまでしか行かないが、測定としてはscatterが一番小さい (系統誤差が最小)
- **StackObject** (stack画像で検出・測光) は MeanObject よりscatterが大きい が、深い撮像 (stack 画像のcomplex PSF shapeによる系統誤差)
- **ForcedMeanObject** は深く, scatterも 比較的小さい


明るめの天体はMeanObject, 暗めの天体はForcedMeanObjectを使う.



Galaxy magnitudes

$g-r$ vs. r color-magnitude diagram

stars are indicated in blue and galaxies in red.



Date product: Photometry Catalogues

Pan-STARRS Catalog Search

We want your feedback on MAST catalogs! Please take five minutes to fill out [this survey](#)! Thank you.

The Panoramic Survey Telescope & Rapid Response System (Pan-STARRS or PS1) is a wide-field imaging facility developed at the University of Hawaii's Institute for Astronomy for a variety of scientific studies from the nearby to the very distant Universe. The PS1 catalog includes measurements in five filters (grizy) covering 30,000 square degrees of the sky north of declination -30 degrees, with typically ~12 epochs for each filter. This interface allows searches for the mean measurements and the deeper stacked measurements from images combining all the epochs. The DR2 release also includes the detection catalog containing all the multi-epoch observations.

Target Supply the central coordinates or target name.

Coordinates

Target name

Target

Mrk590

Resolve Target

Search Radius

1

Arcminutes

Max = 30 arcminutes

Crossmatch a List of Targets Upload a CSV file.

What to Search Select the catalog type and release to search.

Release

PS1 DR1

PS1 DR2

Catalog

Mean object

Stacked object

Forced mean
object

Detections

[View Forced Mean Object Table](#)

Display Columns Select the columns that will be displayed. 58 selected

Search Conditions Select the rows that will be displayed.

+ Add condition

raMean

is less than or equal to

360

Remove

Search Catalog

MASTでホストされている

[Mikulski Archive for Space Telescopes](#)

web interfaceで 直感的に
測光カタログデータを取得可能

4つの必要十分なcatalogs
のみのsimpleなinterface.
よりカスタマイズした
catalog accessは
Casjobs SQL(後述)を用いる.

詳細は → <https://catalogs.mast.stsci.edu/docs/panstarrs.html>
<https://catalogs.mast.stsci.edu/panstarrs/>

Date product: Photometry Catalogues

例: ForcedMean

dr2/forced_mean Target: mrk590/NGC 0863 RA: 33.63984 Dec: -0.76669 Radius: 0.08333°													↓ Export
objName	objID	nDetections	gnTotal	gFPSFFlux	gFPSFFluxErr	gFPSFFluxStd	gFKronFlux	gFKronFluxErr	gFKronFluxStd	gFApFlux	gFApFluxErr	gFApFluxStd	
PSO J033.5587-00.7824	107060335586721562	0	12	0.000005807060006191023	0.000002623409955049283	7.042540073394775	0.000025748700863914564	0.000007156500032579061	11.137200355529785	0.000022776499463361688	0.00000798643031885149	66.352096557	
PSO J033.5589-00.7745	107070335588340943	1	12	0.000002539220076869242	5.500199904417968e-7	1.3470300436019897	0.000004123330199945485	0.0000012260200037417235	2.291029930114746	0.000003544389983289875	0.00000108348001504055	3.5850400924	
PSO J033.5591-00.7735	107070335590502293	1	12	0.0000010838200523721753	3.6332801300886786e-7	0.7818589806556702	0.0000032512100460735383	0.0000017942900285561336	1.2332799434661865	0.000002747430016825092	4.129690012177889e-7	1.2400100231	
PSO J033.5595-00.7692	107070335594837354	0	12	6.295559842328657e-7	3.3027100698745926e-7	0.8068040013313293	7.933979873087083e-7	6.634990086240578e-7	0.6189829707145691	9.819150363910012e-7	6.75969999747681e-7	1.6883599758	
PSO J033.5599-00.7753	107060335600719805	1	12	0.0000028389399631123524	2.8340699032014527e-7	0.5684890151023865	0.00000544121985512902	0.0000016168499996638275	3.846980094909668	0.000004134909886488458	0.0000012603500181285199	6.1825699806	
PSO J033.5600-00.7609	107080335599857300	0	12	0.0000019649698970169993	0.0000016182100353034912	2.4534499645233154	-5.802950227007386e-7	0.0000060521701925608795	6.520579814910889	-3.4080900945809844e-7	0.0000037224399420665577	12.996199607	
PSO J033.5612-00.7398	107110335612292683	10	12	0.0000016931500113059883	5.44421027370845e-7	0.7249739766120911	0.000004125310169911245	9.528529858471302e-7	0.4880790114402771	0.0000023955199139891192	0.0000014977099453972187	5.8249998092	
PSO J033.5615-00.7816	107060335614272476	37	12	0.000008689099558978342	6.617070198444708e-7	1.096869945526123	0.000010779300282592885	0.0000015390500038847676	2.8405098915100098	0.000012925200280733407	0.000001390450051985681	5.7856302261	
PSO J033.5618-00.7801	107060335618184276	0	12	0.0000012176400332464254	4.620069944394345e-7	0.4257600009441376	0.000006922030024725245	0.0000018918200339612667	5.932799816131592	0.0000057099400692095514	6.137299806141527e-7	0.4445729851	
PSO J033.5620-00.7930	107040335619438541	0	26	0.0000014703100532642566	1.8340099927627307e-7	0.24372799694538116	0.000005886419785383623	0.000002498989942978369	3.867919921875	3.322359987123491e-7	3.0067099032748956e-7	0.57773149728	
PSO J033.5621-00.7694	107070335620327211	4	12	0.0000016579999737587059	8.167339728970546e-7	1.2163699865341187	0.000003966769781982293	2.283759954480047e-7	0.26725199818611145	0.000002125000037267455	0.0000016852000044309534	4.8672299385	
PSO J033.5627-00.7827	107060335626901243	9	12	0.0000014293899539552513	5.623360266326927e-7	0.5692870020866394	0.00000632859018878662	0.000006251209924812429	9.926799774169922	0.0000017160100469482131	0.000004520959919318557	43.058601379	
PSO J033.5638-00.7630	107080335637454773	0	12	0.000007725209798081778	0.000001791819954632956	133.08799743652344	0.00004299830106901936	0.00003606699829106219	275.9549865722656	0.00003117090091109276	0.000029566999728558585	1769.3199462	
PSO J033.5639-00.7847	107050335639048834	0	26	0.0000025740400815266185	2.5466201236667985e-7	0.27830201387405396	0.0000018729399471340002	0.0000010906300076385378	3.0968499183654785	0.0000033441199320805026	0.000001358579993393505	10.532999992	
PSO J033.5640-00.7615	107080335639906580	0	12	0.000008874900231603533	0.0000014108500181464478	-999	0.00003671320155262947	0.0000027677199341269443	-999	0.00003091320104431361	0.0000014108500181464478	-999	

PSF flux, Kron flux, aperture flux … などが利用できる。

↑ Fluxは *Jansky* unit

* ForcedWarp, ForcedMean [FPSFFlux, FKronFlux など]はマイナス符号の場合もある

The Pan-STARRS1 data archive home page



Welcome to the starting point for access to data from the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS). This page provides a brief summary of the facilities and data products to guide Pan-STARRS archive users. More complete information is provided on linked pages (see below).

[Pan-STARRS](#) is a system for wide-field astronomical imaging developed and operated by the Institute for Astronomy at the University of Hawaii. Pan-STARRS1 (PS1) is the first part of Pan-STARRS to be completed and is the basis for both Data Releases 1 and 2 (DR1 and DR2). The PS1 survey used a 1.8 meter telescope and its 1.4 Gigapixel camera (GPC1; see [PS1 GPC1 camera](#)) to image the sky in five broadband [filters](#) (g, r, i, z, y). The PS1 Science Consortium funded the operation of the Pan-STARRS1 telescope, situated at [Haleakala Observatories](#) near the summit of Haleakala in Hawaii, for the purposes of astronomical research. The PS1 consortium is made up of astronomers and engineers from [14 institutions from six countries](#).

The data from PS1 are archived at the [Space Telescope Science Institute](#) (STScI) in Baltimore Maryland, and can be accessed through MAST, the Mikulski Archive for Space Telescopes. Additional support for the PS1 public science archive is provided by the [Gordon and Betty Moore Foundation](#).

The [web site for Pan-STARRS1 in Hawaii](#) also describes the project.

Quick links to the MAST PS1 Archive User Interface

The current PS1 data release is DR2 (2019 January 28). Use the following links to jump right to the MAST PS1 Science Archive interfaces and get started using PS1 data!

Object Catalog Search: <https://catalogs.mast.stsci.edu/>

Image Cutout Server: <https://ps1images.stsci.edu/cgi-bin/ps1cutouts>

Casjobs SQL Server: <https://mastweb.stsci.edu/ps1casjobs>

See [How to retrieve and use PS1 data](#) for more information on these access mechanisms. Note that all of these allow programmatic scripted access, and there are sample Python Jupyter notebook scripts.

The following [image](#) and [catalog](#) products are available in **DR2**:

- [Warp images](#) are the result of resampling and realigning the camera images into a skycell of the [PS1 Sky Tessellation](#), a set of common pixel-grid images with simple projections from the sky. Warps are astrometrically and photometrically calibrated.
- Extracted photometry for point sources and extended objects from the [warp images](#) are available in DR2. The DR2 source database allows users to extract information on any time variation of source photometry or astrometry.
- [Stack images](#) are co-added images made from the multiple exposures taken over the survey. Stacks provide the best signal-to-noise, and the source catalogs created from the stacks are recommended as a starting point for data analysis.
- [PS1 source extraction and catalogs](#) describes how point-source and extended-object photometry is measured from the [stack images](#). These source extractions are made available in DR1.
- Mean values of the point-source and extended-object photometry from the [warp images](#) are also available in DR1.

PS1 user information topics

There are five primary informational topics, with links below, that provide further details:

- [The Pan-STARRS1 facilities and hardware](#)
- [PS1 Data processing procedures](#)
- [PS1 Image data products](#)
- [PS1 Source extraction and catalogs](#)
- [How to retrieve and use PS1 data](#)
- [PS1 sample queries](#)

Detailed explanations of the products,
sample Python Jupyter notebooks, ...


<https://panstarrs.stsci.edu/>

Data accessの例 (image):


image cutout web interfaceにpythonからaccess

例: cutout stack fits画像をdownload & plot

```
1
2 # Mrk590 position
3 ra = 033.6398378956147
4 dec = -00.7667167022724
5 size = 1280
6
7 #####
8 # accessするurlを生成
9 #####
10 fitsurl = geturl(ra, dec, size=size, filters="i", format="fits")
11 # --> 'https://ps1images.stsci.edu/cgi-bin/fitscut.cgi?ra=33.6398378956&dec=
12 #      -0.766716702272&size=1280&format=fits&red=/rings.v3.skycell/1240/080/
13 #      rings.v3.skycell.1240.080.stk.i.unconv.fits'
14
15 #####
16 # 画像をlocalに読み込む (HDU)
17 #####
18
19 image_url = fitsurl[0]
20 variance_url = fitsurl[0].replace("unconv.", "unconv.wt.")
21
22 fh = fits.open(image_url)
23 fh_variance = fits.open(variance_url)
24
25 #####
26 # Fits画像をlocal fileに保存する.
27 #####
28
29 out_image = fh[0].data
30 out_variance = fh_variance[0].data
31
32 os.system("rm " + out_image)
33 os.system("rm " + out_variance)
34 fh.writeto(out_image)
35 fh_variance.writeto(out_variance)
36
37
```



Stack



Stack variance (wt)

→ python/photutilsなどで測光解析できる (stack画像はゼロ点等級=25等; fits headerをみよ)


上記で用いたpython tasksは <https://ps1images.stsci.edu/ps1image.html> にある.

<https://outerspace.stsci.edu/display/PANSTARRS/How+to+retrieve+and+use+PS1+data>


https://ps1images.stsci.edu/ps1_dr2_api.html

Data accessの例 (image): image cutout web interfaceにpythonからaccess

例: cutout stack fits画像をdownload & plot



Stack



Stack variance (wt)

`rings.v3.skycell.1240.080.stk.i.unconv.fits`

“rings.v3で定義されるx=1240,y=080のskycellのstack画像, i-band”

Data accessの例 (image):

image cutout web interfaceにpythonからaccess


例: cutout stack fits画像をdownload & plot

普通のfits imageなので、ds9で表示したり


```
ds9 -zscale \  
rings.v3.skycell.1240.080.stk.i.unconv.fits \  
rings.v3.skycell.1240.080.stk.i.unconv.wt.fits
```

astropy.ioでpythonで読み込んだりできる

```
python  
from astropy.io import fits  
hdu = fits.open("rings.v3.skycell.1240.080.stk.i.unconv.fits")  
hdu[0].header['EXPTIME']  
hdu[0].header['HIERARCH FPA.ZP']
```



Stack



Stack variance (wt)

ds9などで、stack画像の分散と
variance mapの値が大体あっているかを
checkしてみると、安心して使える

Data accessの例 (catalog):

MAST web API にpythonからaccess

- MASTのAPIにqueryを送り,
`python.requests.get`
でpythonに取り込む

```
1 #
2 # This searches the mean object catalog for objects within 50 arcsec of Mrk590.
3 # Note that the results are restricted to objects with nDetections>1, where nDetections is the total number
4 # of times the object was detected on the single-epoch images in any filter at any time.
5 # Objects with nDetections=1 tend to be artifacts, so this is a quick way to eliminate most spurious objects
6 # from the catalog.
7 #
8 # Mrk590の周辺r=50arcsecのMeanObjectを検索.
9 #
10 # nDetections>1 (どのfilterでもよいので, 2回以上single-epoch画像で検出されている)という条件をつける.
11 # nDetections=1は,ほとんどの場合spurious objects(ゴミ)なので, nDetections>1の条件でかなりの割合のゴミを排除できる.
12 #
13 # "Mean" catalogのcolumnの説明は
14 # https://outerspace.stsci.edu/display/PANSTARRS/PS1+MeanObjectView+table+fields
15 #
16
17
18 ra      = 33.6398378956147
19 dec     = -0.7667167022724
20 radius  = 50.0/3600.0
21 constraints = {'nDetections.gt':1}
22
23 # Mean catalogの中のいくつかのcolumnを抽出する.
24 columns = """objID,raMean,decMean,nDetections,ng,nr,ni,nz,ny,
25 gMeanPSFMag,rMeanPSFMag,iMeanPSFMag,zMeanPSFMag""".split(',')
26 columns = [x.strip() for x in columns]
27 columns = [x for x in columns if x and not x.startswith('#')]
28
29 # ps1cone: web APIにquery urlを送り、APIが出力したcsvをresultsに読み込む.
30 results = ps1cone(ra,dec,radius,release='dr2',columns=columns,verbose=True,**constraints)
31
32 ## print first few lines
33 #lines = results.split('\n')
34 #print(len(lines),"rows in results -- first 5 rows:")
35 #print('\n'.join(lines[:6]))
36
37 # astropy.table形式に変換.
38 tab = ascii.read(results)
39 # improve the format
40 for filter in 'grizy':
41     col = filter+'MeanPSFMag'
42     try:
43         tab[col].format = ".4f"
44         tab[col][tab[col] == -999.0] = np.nan
45     except KeyError:
46         print("{} not found".format(col))
47
48 # 表示
49 tab
50
```

<https://catalogs.mast.stsci.edu/api/v0.1/panstarrs/dr2/mean.csv?dec=-0.766716702272&radius=0.0138888888889&ra=33.6398378956&columns=%5BobjID%2CraMean%2CdecMean%2CnDetections%2Cng%2Cnr%2Cni%2Cnz%2Cny%2CgMeanPSFMag%2CrMeanPSFMag%2CiMeanPSFMag%2CzMeanPSFMag%2CyMeanPSFMag%2CSD%2CnDetections.gt=1>

<Table length=31>


objID	raMean	decMean	nDetections	ng	nr	ni	nz	ny	gMeanPSFMag	rMeanPSFMag	iMeanPSFMag	zMeanPSFMag
int64	float64	float64	int64	int64	int64	int64	int64	int64	float64	float64	float64	float64
107060336410857640	33.64110938	-0.77741577	2	0	1	1	0	0	nan	21.4057	21.1266	nan
107060336422858569	33.64226204	-0.77667082	2	0	0	1	1	0	nan	nan	21.4750	20.7051
107070336320732430	33.63212833	-0.77334054	2	0	1	1	0	0	nan	21.3538	21.6887	nan
107070336336271423	33.63357657	-0.77408146	2	2	0	0	0	0	22.6118	nan	nan	nan
107070336390957480	33.63909535	-0.76912359	2	0	2	0	0	0	nan	nan	nan	nan
107070336394414750	33.6393077	-0.77123597	3	1	0	2	0	0	20.4766	nan	19.3916	nan
107070336402931774	33.64034712	-0.77376801	2	0	0	0	1	1	nan	nan	nan	20.6208
107070336426286779	33.64280825	-0.76983767	2	1	0	1	0	0	19.6913	nan	19.9995	nan
107070336454475298	33.64545022	-0.77098068	8	0	5	3	0	0	nan	20.3723	20.4459	nan
107080336281025128	33.62812549	-0.76267443	2	2	0	0	0	0	21.5735	nan	nan	nan

<https://catalogs.mast.stsci.edu/docs/panstarrs.html>

scriptは次のURLから持ってきた: https://ps1images.stsci.edu/ps1_dr2_api.html

Data accessの例 (catalog):

MAST web API にpythonからaccess



- MASTのAPIにqueryを送り,
`python.requests.get`
でpythonに取り込む

← *Detection* tableの
PSF flux測定値 light curve
(0.4863日周期で折りたたみ)


星 (点源)の測光light curveは
素性が良い

<https://catalogs.mast.stsci.edu/docs/panstarrs.html>

scriptは次のURLから持ってきた: https://ps1images.stsci.edu/ps1_dr2_api.html

Data accessの例 (catalog):

MAST web API にpythonからaccess



- MASTのAPIにqueryを送り,
`python.requests.get`
でpythonに取り込む

← *Detection* tableの
Kron flux測定値 light curve

AGN (広がった天体)の測光light curve
は要注意


<https://catalogs.mast.stsci.edu/docs/panstarrs.html>

scriptは次のURLから持ってきた: https://ps1images.stsci.edu/ps1_dr2_api.html

Data accessの例 (catalog): Casjobs SQL interface

- 大量のカatalogデータを読み込んだり、Catalog同士を組み合わせたりする場合、SQL serverにQueryを投げる
 - Create a MAST Query / casjobs account
 - Write a SQL query to get desired SQL table
 - Submit the query

SQLサーバで利用できるテーブル一覧→




The screenshot shows the MAST Query / CasJobs web interface. At the top, there is a logo and the title "MAST Query / CasJobs". Below the title is a navigation bar with links: Home, Help, Tools, Query, History, and MyDB. A dropdown menu is set to "PanSTARRS_DR2". The main content area is divided into sections: Views, Tables, Functions, and Procedures. Below these sections, there is a table listing available tables and views. The table has three columns: Rows, kB, and Name. The list includes various astronomical data tables and views, such as Detection, DetectionObjectView, DiffDetection, DiffDetObjectView, ForcedGalaxyModelView, ForcedGalaxyShape, ForcedMeanLensing, ForcedMeanObjectView, ForcedWarpExtended, ForcedWarpLensing, ForcedWarpMasked, ForcedWarpMeasurement, GaiaDR2_PS1DR2_2arcsec, MeanObject, MeanObjectView, StackApFlx, StackApFlxExGalCon6, StackApFlxExGalCon6ObjectView, StackApFlxExGalCon8, StackApFlxExGalCon8ObjectView, StackApFlxExGalUnc, StackApFlxExGalUncObjectView, StackApFlxObjectView, StackModelFitDeV, StackModelFitDeVObjectView, StackModelFitExp, StackModelFitExpObjectView, StackModelFitExtra, StackModelFitPetObjectView, StackModelFitSer, StackModelFitSerObjectView, StackModelObjectView, StackObjectAttributes, StackObjectView, StackPetrosian, and StackToImage.

Rows	kB	Name
NA	NA	Detection
NA	NA	DetectionObjectView
NA	NA	DiffDetection
NA	NA	DiffDetObjectView
NA	NA	ForcedGalaxyModelView
NA	NA	ForcedGalaxyShape
NA	NA	ForcedMeanLensing
NA	NA	ForcedMeanObjectView
NA	NA	ForcedWarpExtended
NA	NA	ForcedWarpLensing
NA	NA	ForcedWarpMasked
NA	NA	ForcedWarpMeasurement
NA	NA	GaiaDR2_PS1DR2_2arcsec
NA	NA	MeanObject
NA	NA	MeanObjectView
NA	NA	StackApFlx
NA	NA	StackApFlxExGalCon6
NA	NA	StackApFlxExGalCon6ObjectView
NA	NA	StackApFlxExGalCon8
NA	NA	StackApFlxExGalCon8ObjectView
NA	NA	StackApFlxExGalUnc
NA	NA	StackApFlxExGalUncObjectView
NA	NA	StackApFlxObjectView
NA	NA	StackModelFitDeV
NA	NA	StackModelFitDeVObjectView
NA	NA	StackModelFitExp
NA	NA	StackModelFitExpObjectView
NA	NA	StackModelFitExtra
NA	NA	StackModelFitPetObjectView
NA	NA	StackModelFitSer
NA	NA	StackModelFitSerObjectView
NA	NA	StackModelObjectView
NA	NA	StackObjectAttributes
NA	NA	StackObjectView
NA	NA	StackPetrosian
NA	NA	StackToImage

Data accessの例 (catalog):

Casjobs SQL interface



The screenshot shows the MAST Query / CasJobs web interface. At the top is a navigation bar with links: Home, Help, Tools, Query, History, MyDB, Import, Groups, Output, Profile, Queues, and Logout. Below the navigation bar is a header section with three tabs: Context, Table (optional), and Task Name. The Context tab is active, showing a dropdown menu with 'PanSTARRS_DR2' selected. The Table (optional) tab shows 'MyTable' and the Task Name tab shows 'My Query'. Below the tabs is a section with buttons: Samples, Recent, and Clear, followed by 'Line 1, Col 1'. The main area displays a sample SQL query for a rectangle search for PS1 objects. A red arrow points from the Japanese text at the bottom to the first line of the SQL query.

MAST Query / CasJobs

Home Help Tools Query History MyDB Import Groups Output Profile Queues Logout

Context Table (optional) Task Name

PanSTARRS_DR2 ▼ MyTable My Query

Samples Recent Clear Line 1, Col 1

```
-- Rectangle search for PS1 objects within a rectangular patch of sky
-- Restrict to nDetections>1 to reduce spurious objects
-- Returns PSF magnitudes and errors in grizy

select o.objID, o.nDetections, o.raMean, o.decMean,
       m.gMeanPSFMag, m.gMeanPSFMagErr, m.rMeanPSFMag, m.rMeanPSFMagErr,
       m.iMeanPSFMag, m.iMeanPSFMagErr, m.zMeanPSFMag, m.zMeanPSFMagErr,
       m.yMeanPSFMag, m.yMeanPSFMagErr
from ObjectThin o
inner join MeanObject m on o.objID=m.objID
where
  o.raMean between 100.0 and 100.1
and o.decMean between 0.0 and 0.1
and o.nDetections>1
```

Sample SQL queryも充実
(SDSS SQLと同じ)

Data accessの例 (catalog):

Casjobs SQL interface

PS1 Sample queries

Armin Rest が作成し、2 13, 2019 に Rick White が最終更新


On this page

- [Get high-fidelity stars in given area](#)
 - [Goal/Description](#)
 - [Tables used](#)
 - [Query](#)
 - [Results](#)
- [Galaxy Candidates for K2 SN Search](#)
 - [Goal/Description](#)
 - [Query](#)
 - [Results](#)
- [Obtain lightcurves for a given set of objects](#)
 - [Goal/Description](#)
 - [Query #1: Match Catalina RR Lyrae stars to PS1 objects](#)
 - [Results \(25 minutes\)](#)
 - [Query #2: Get the detections](#)
 - [Results \(30 seconds\)](#)
 - [Query #3: Get the forced detections](#)
 - [Results \(3:45 minutes\)](#)
- [Obtain lightcurves for a single object](#)
 - [Goal/Description](#)
 - [Query #1: Get the ObjID for the star](#)
 - [Results \(1 entry, < 1 min\)](#)
 - [Query #2: Get the detections](#)
 - [Results \(92 entries, < 1 min\)](#)
 - [Query #3: Get the forced detections](#)
 - [Results \(84 entries, <1 min\)](#)
- [Select mean objects and associated detections within 3 deg of HDF; compare photometry and astrometry with Gaia DR2](#)
 - [Step 1: Obtain mean objects in cone search, filtering for objects with a high probability of Gaia matches](#)
 - [Step 2: Obtain detections associated with objects selected in Step 1](#)
 - [Step 3: Match objects and detections with the closest Gaia DR2 source](#)
 - [Step 4: compare Gaia and PanSTARRS magnitudes for point sources](#)
 - [Step 5: Compute mean detection positions and compare to Gaia astrometry](#)

Data accessの例 (catalog):

Casjobs SQL interface

CSS J030521.9+013231 (変光星RR Lyrae)




```
script3 = "\
SELECT \
o.objID, o.raMean, o.decMean, \
fwm.detectID, \
fwm.obstime, fwm.exptime, fwm.airmass, fwm.Fpsfflux, fwm.FpsffluxErr, \
fwm.FpsfQf, fwm.FpsfQfPerfect, fwm.FpsfChiSq, fwm.zp, fwm.FapFlux, \
fwm.FapFluxErr, \
fwm.forcedWarpID, fwm.filterID, \
fwm.Fsky, fwm.Fskyerr, fwm.Finfoflag, fwm.Finfoflag2, fwm.Finfoflag3, \
o.qualityFlag, \
o.gMeanPSFMag, o.gMeanPSFMagErr, o.gMeanPSFMagNpt, \
o.rMeanPSFMag, o.rMeanPSFMagErr, o.rMeanPSFMagNpt, \
o.iMeanPSFMag, o.iMeanPSFMagErr, o.iMeanPSFMagNpt, \
o.zMeanPSFMag, o.zMeanPSFMagErr, o.zMeanPSFMagNpt, \
o.yMeanPSFMag, o.yMeanPSFMagErr, o.yMeanPSFMagNpt, \
o.rMeanKronMag, o.rMeanKronMagErr, \
o.nDetections, o.ng, o.nr, o.ni, o.nz, o.ny, \
o.gFlags, o.gQfPerfect, \
o.rFlags, o.rQfPerfect, \
o.iFlags, o.iQfPerfect, \
o.zFlags, o.zQfPerfect, \
o.yFlags, o.yQfPerfect \
INTO mydb.[RRL_584630948352256_PS1forceddet] \
FROM mydb.[RRL_584630948352256_PS1] o \
JOIN ForcedWarpMeasurements fwm on fwm.ObjID = o.ObjID \
"
"java -jar casjobs.jar run -t 'PanSTARRS_DR2' -n 'RRL_584630948352256_PS1forceddet' " + "'" + script3 + "
```

↑ Command line tool casjobsで
CasjobsにSQL queryをsubmit
→ .csvでdownload


← *ForcedWarpMeasurements* tableの
PSF flux測定値 light curve
(0.55547日周期で折りたたみ)



SkyMapper Southern Survey (SMSS)




<https://sciencesprings.wordpress.com/2020/10/10/meet-the-siding-spring-observatory/>




SkyMapper Southern Survey (SMSS)

- 2014 — ongoing. 南天 2π steradian.
- 最終目標は 20 — 22 mag in u, v, g, r, i, z -band


u -bandが2つ (QSO; 金属欠乏星selection)



1.35m口径, 5.7deg²



32 E2V CCD4482 devices: 2048x4096 15 micron pixel CCDs



Bessell et al. (2011).

Shallow survey & Main Survey

Shallow Survey

Wolf+2018

Filter	$\lambda_{\text{cen}}/\Delta\lambda$ (nm)	R_{band}	t_{exp} (s)	N_{image}	10 σ -limit (ABmag)	FWHM (arcsec)
<i>u</i>	349/42	4.294	40	10909	17.9	3.1
<i>v</i>	384/28	4.026	20	10779	17.7	2.9
<i>g</i>	510/156	2.986	5	11872	18.0	2.6
<i>r</i>	617/156	2.288	5	11515	18.0	2.4
<i>i</i>	779/140	1.588	10	10698	18.0	2.3
<i>z</i>	916/84	1.206	20	11067	18.0	2.3

Main Survey

Onken+2020

5 σ depth (mag.)	
<i>u</i>	20
<i>v</i>	21
<i>g</i>	22
<i>r</i>	22
<i>i</i>	21
<i>z</i>	20

- **Shallow Survey:** 短いexposureで 2π ステラジアンを掃き、~18等mapを作る. 2014-2015で完了. Data Release 1で公開.
- **Main Survey:** Shallow Survey完了後, ~22等を目指して長めのexposureで 2π ステラジアンを掃く. 現在進行中. Data Release 2で部分的に公開.
 - DR2の段階ではsingle-epochの画像, 測光値やその平均値のみ (instantaneous photometry only). stack画像やstack photometryは公開されていない ... 現状だとあまり使い勝手がよくない
 - Stack画像の公開はDR3から (Onken+2020).

- Home
- About SkyMapper
- Purchase Time**
- Cite & Acknowledge
- Latest News
- Survey Information
- Data Releases**
- Survey Progress**
- View Publications**
- Publication Statistics**
- Data Access
- How To Access
- Object Viewer
- Cone Search
- Image Cutouts
- Full Catalogue Search
- Table Metadata
- Sky Viewer
- Protected Science

Using a specially-built, 1.3-meter telescope at Siding Spring Observatory near Coonabarabran, the **SkyMapper Southern Survey** is producing a high-fidelity digital record of the entire southern sky for Australian astronomers.

[Learn More →](#)



SkyMapper's *Southern Survey* is led by the **Research School of Astronomy and Astrophysics** at the **Australian National University**, in collaboration with seven Australian universities and the Australian Astronomical Observatory. The goal of the project is to create a deep, multi-epoch, multi-colour digital survey of the entire southern sky. This will facilitate a broad range of exciting science, including discovering the oldest stars in the Galaxy, finding new dwarf galaxies in orbit around the Milky Way, and measuring the effects of Dark Energy on the Universe through nearby supernovae.

On this site you can **read about SkyMapper** and its **surveys**, browse the latest **data releases** and query **images and catalogues** using simple forms or Virtual Observatory web services. Before downloading or publishing data we ask that you review the **policies** section to familiarise yourself with the authorship and protected science policies. As part of SkyMapper's status as a National Facility, Australian

Latest Data Release

DR2

Feb 27, 2019

21,000 deg²
 505,176,667 objects
 121,494 exposures
 4.7 billion detections
 Matched against 2MASS,
 AllWISE, ATLAS Refcat2,
 Gaia, GALEX, Pan-STARRS1,
 and UCAC4


[Learn More →](#)

Follow [@SkyMapper](#) on Twitter



SkyMapper Telescope
 @SkyMapper

Data Release 2 from the SkyMapper Southern Survey is



SkyViewer

Southern Sky Viewer

This viewer was constructed using the Hierarchical Progressive Survey format (**HiPS**) used by **Aladin-Lite**. Navigate and zoom in/out using your mouse or the controls provided. You can search for coordinates or objects and change the displayed image layer to show other all-sky surveys using the icons in the top left, and maximise the viewer using the arrows in the top right.



“見る”だけならSkyViewerで十分
(stack画像ではなく, おそらく best single-epoch画像)

Object catalog

dr2.master
dr2.ccds
dr2.images
dr2.mosaic
dr2.photometry

- DR2測光カタログは2種類
 - dr2.master: 複数epochの測光値の平均値
[PanSTARRSのMeanObjectのようなもの]
 - dr2.photometry: 各epochの測光値
[PanSTARRSのDetectionsのようなもの]

Cone Search Service (dr2.masterのみのシンプルなインターフェイス)

This service allows you to query the **master** table of mean photometry and object parameters around a given position. Please specify either Equatorial/Galactic coordinates or a SIMBAD-resolvable object name as well as a search radius in arcseconds (maximum 3,600" or 2,000 rows). You may increase the number of columns returned by increasing the metadata verbosity parameter. This service is also accessible through VO-aware clients like **TOPCAT** using the IVOA Cone Search standard. See the **How To Access** page for more information.

Note: By using SkyMapper data, you agree to observe the **SkyMapper policies** published on this website.

▶ Data Release

▶ Search Parameters

Please input either a set of coordinates **or** a SIMBAD name.

Coordinate system:

Coordinates:

Separate longitude & latitude with space(s)

Valid coordinate types, e.g.:

4h30m15.82s

OR

Target name:

A SIMBAD-compatible target name

Search radius:

Arcseconds

Metadata verbosity:

Detail in returned metadata

<http://skymapper.anu.edu.au/cone-search/>

Object catalog


dr2.master
dr2.ccds
dr2.images
dr2.mosaic
dr2.photometry

- DR2測光カタログは2種類
 - dr2.master: 複数epochの測光値の平均値
[PanSTARRSのMeanObjectのようなもの]
 - dr2.photometry: 各epochの測光値
[PanSTARRSのDetectionsのようなもの]
- dr2.masterカタログのデータを、あるRA, Decl周辺で取得し, XML形式でoutput:
<http://skymapper.anu.edu.au/sm-cone/public/query?RA=33.639837321&DEC=-0.766719986&SR=0.1&VERB=3&CATALOG=dr2.master&RESPONSEFORMAT=XML>
- dr2.photometryデータをCSV形式でoutput
<http://skymapper.anu.edu.au/sm-cone/public/query?RA=33.639837321&DEC=-0.766719986&SR=0.1&VERB=3&CATALOG=dr2.photometry&RESPONSEFORMAT=CSV>

*python*からWeb-API Image cutout service → download

```
: # SkyMapper image cone search CSV pandas DataFrame .  
#  
# query Mrk590 CCD .  
#  
# http://api.skymapper.nci.org.au/public/siap/dr2/query?\  
# POS=33.639837321,-0.766719986&SIZE=0.17,0.17&FORMAT=image/  
  ↪fits&INTERSECT=center&%20&RESPONSEFORMAT=CSV  
  
url = "http://api.skymapper.nci.org.au/public/siap/dr2/query?POS=33.  
  ↪639837321,-0.766719986&SIZE=0.17,0.17&FORMAT=image/  
  ↪fits&INTERSECT=center&%20&RESPONSEFORMAT=CSV"  
  
#r = requests.get(url, params=payload)  
r = requests.get(url)  
  
df = pd.read_csv(io.BytesIO(r.content),sep=",")
```

		image_name	format	band	\
0	SkyMapper_u_20140808191436-29_33.623-0.767_489...	image/fits	u		
1	SkyMapper_v_20140808191542-29_33.623-0.767_488...	image/fits	v		
2	SkyMapper_g_20140808191626-29_33.623-0.767_489...	image/fits	g		



PanSTARRSの場合の利用法と同様

*python*からWeb-API catalog cone search → download & plot

```
# dr2.photometry CSV output
# Mrk590 (SR=search radius in deg unit.)
# http://skymapper.anu.edu.au/sm-cone/public/query?RA=33.639837321&DEC=-0.
  ↳766719986&SR=0.0003&VERB=3&CATALOG=dr2.photometry&RESPONSEFORMAT=CSV
#


url = "http://skymapper.anu.edu.au/sm-cone/public/query?RA=33.639837321&DEC=-0.
  ↳766719986&SR=0.0003&VERB=3&CATALOG=dr2.photometry&RESPONSEFORMAT=CSV"

#r = requests.get(url, params=payload)
r = requests.get(url)

df_phot = pd.read_csv(io.BytesIO(r.content), sep=",")
```


```
df_phot.columns
```

```
Index([u'r_cntr', u'object_id', u'object_id_local', u'image_id', u'ccd',
      u'filter', u'ra_img', u'decl_img', u'x_img', u'y_img', u'flags',
      u'nimaflags', u'imaflags', u'background', u'flux_max', u'mu_max',
      u'class_star', u'a', u'e_a', u'b', u'e_b', u'pa', u'e_pa', u'elong',
      u'fwhm', u'radius_petro', u'radius_kron', u'radius_frac20',
      u'radius_frac50', u'radius_frac90', u'chi2_psf', u'flux_psf',
      u'e_flux_psf', u'mag_psf', u'e_mag_psf', u'flux_petro', u'e_flux_petro',
      u'mag_petro', u'e_mag_petro', u'flux_kron', u'e_flux_kron', u'mag_kron',
```



Subaru/HSC public/internal data release

- SDSSのデータベース同等 … sky viewer, SQL catalog/data archive server など共通



<https://hsc-release.mtk.nao.ac.jp/hscMap-pdr2/app/>

← → ↻ hscdata.mtk.nao.ac.jp:4443/datasearch/

about schema browser list jobs summary table search direct SQL search misc hscMap (s19a) (s18a) (s17a)

name:

release:

SQL: Press ctrl-enter to preview

```
1 -- Simple area search based on the range of RA and DEC getting all object
2 -- with i band cmodel magnitudes smaller than 25.5, and in the area in
3 -- the range RA J2000 between 34.0 and 36.0 degrees and DEC J2000 -5.0
4 -- and -4.5 degrees.
5
6 -- WARNING:
7 -- --> Remove 'LIMIT 10' for your query
8 -- --> Edit the schema name 's20a_wide' for your query.
9
10 SELECT
11     object_id
12     ,ra
13     ,dec
14     ,i_cmodel_mag
15     ,i_cmodel_magerr
16     ,y_cmodel_mag
17     ,y_cmodel_magerr
18     ,i_cmodel_mag - y_cmodel_mag AS i_y
19 FROM
20     s20a_wide.forced
21 WHERE
22     boxSearch(coord, 34.0, 36.0, -5.0, -4.5)
23     /* is equivalent to
24        ra BETWEEN 34.0 AND 36.0
25        AND dec BETWEEN -5.0 AND -4.5
26        but boxSearch() is much faster
27     */
28     AND i_cmodel_mag < 25.5
29     AND isprimary
30 LIMIT 10
31 ;
32
```

example:

- ☐ include SQL in CSV
- ☒ syntax check before enqueueing
If you encounter an error "ERROR: canceling statement due to statement timeout" on enqueueing, please uncheck this.
- ☐ csv ☒ csv.gz ☐ sqlite3 ☐ fits ☐ fast-fits (experimental)

show the estimated query time and the query plan. [What is this?](#) (experimental)

(timeout: 10 seconds) (timeout: 12 hours / output limit: 20 GB)

<https://hsc.mtk.nao.ac.jp/ssp/japanese-data-access/>

Summary

- SDSSレガシーサーベイでカバーされなかった天域に対する撮像サーベイが完了(PanSTARRS 3 π)・進行中(SkyMapper Southern Survey).
他にも, 本スライドでは触れなかった無数の掃天サーベイ, 突発天体サーベイがあり、公開データを提供しているものもある.
- 公開データの構造やアクセス方法はサーベイ間で共通点が多い.
しかし残念ながら各サーベイの”癖”があるので、**Webページのマニュアルを読み込む必要がある**. Webページには大抵, 多くのquery例が掲載されており、少し改変すれば自分のサイエンスにすぐ使える.
- LSSTなどの将来計画でも、公開データは同様の方法で提供されるだろう.